

Boundary terms for eleven-dimensional supergravity and M -theory

Ian G. Moss*

School of Mathematics, University of Newcastle Upon Tyne, NE1 7RU, UK

(Dated: February 1, 2008)

A new action for eleven dimensional supergravity on a manifold with boundary is presented. The action is a possible low energy limit of M -theory. Previous problems with infinite constants in the action are overcome and a new set of boundary conditions relating the behaviour of the supergravity fields to matter fields are obtained. One effect of these boundary conditions is that matter fields generate gravitational torsion.

One of the phenomenologically interesting low energy limits of M -theory corresponds to eleven-dimensional supergravity on a manifold with a ten-dimensional boundary. Horava and Witten have argued that this theory gives the strongly coupled limit of the $E_8 \times E_8$ heterotic string [1, 2]. The gauge and matter fields propagate on the boundary and couple to the bulk supergravity. Previous attempts to construct these couplings have been hampered by the appearance of the square of the Dirac delta function. This paper presents an improved construction which results in a consistent set of interaction terms.

The interaction terms in the model are constructed as an expansion in $\kappa^{2/3}$, where κ is the eleven dimensional gravitational coupling constant. At leading order, the theory is eleven-dimensional supergravity on the background $R^{10} \times S^1/Z_2$. Since the orbifold S^1/Z_2 is identical to an interval I , with the covering space S^1 , the background has a boundary consisting of two timelike ten-dimensional surfaces or branes. One of these contains the visible matter with which we are familiar and the other contains hidden matter. Gravitational forces are transmitted through the interior which is often called the bulk.

The modified theory described here is presented from the point of view of the manifold with a boundary, rather than the covering space. The action allows for the possibility that the boundary may curve, as we would require for descriptions of cosmology. The main innovations appear in the boundary conditions. In the first place there are corrections to the chirality condition on the gravitino, which are to a large extent forced by the requirements of supersymmetry. There are also boundary conditions for the supergravity three-form which play an important role. The resulting theory is supersymmetric at orders κ^0 , $\kappa^{2/3}$ and $\kappa^{4/3}$.

The notation used here is based on [3]. The metric signature is $- + \dots +$. Vector indices are $I, J, \dots$ in the bulk, $A, B, \dots$ on the boundary and N in the (unit) normal direction. The outward going normal has extrinsic curvature K_{AB} . The exterior derivative of an n -form α is $(d\alpha)_{I_1 \dots I_{n+1}} = (n+1)^{-1} \partial_{[I_1} \alpha_{I_2 \dots I_{n+1}]}$ and the wedge product $(\alpha \wedge \beta)_{I_1 \dots I_m} = m C_n \alpha_{[I_1 \dots I_n} \beta_{I_{n+1} \dots I_m]}$ where $m C_n$ is a binomial coefficient. The gamma matrices satisfy $\{\Gamma_I, \Gamma_J\} = 2g_{IJ}$ and $\Gamma^{I \dots K} = \Gamma^{[I} \dots \Gamma^K]$. The spinors are Majorana, and $\bar{\psi} = \psi^T \Gamma^0$.

The action will be presented first, followed by an account of the boundary conditions. The supersymmetry of the action will be discussed at the end. The supergravity part contains the metric g , gravitino ψ_I and three-form C (with field strength $G = dC$) [4]. The usual supergravity action is

$$S_{SG} = \frac{2}{\kappa^2} \int_{\mathcal{M}} d\mu \left(-\frac{1}{2} R - \frac{1}{2} \bar{\psi}_I \Gamma^{IJK} D(\Omega^*)_{J\psi_K} - \frac{1}{48} G_{IJKL} G^{IJKL} \right. \\ \left. - \frac{\sqrt{2}}{192} (\bar{\psi}_I \Gamma^{IJKLMP} \psi_P + 12 \bar{\psi}^J \Gamma^{KL} \psi^M) G_{JKLM} - \frac{\sqrt{2}}{10!} \epsilon^{I_1 \dots I_{11}} (C \wedge G \wedge G)_{I_1 \dots I_{11}} \right) \quad (1)$$

In the 1.5 order formalism used here, the spin connection Ω is varied as an independent field. The combination $\Omega^* = (\Omega + \hat{\Omega})/2$, where hats denote a standardised subtraction of gravitino terms to make a supercovariant expression.

The background for the low energy limit of M theory of interest here contains a manifold with timelike boundaries. The boundary conditions are fixed by symmetries on the covering space. On a manifold with boundary, the Einstein action is supplemented by an extrinsic curvature term [5, 6, 7]. The supersymmetric version has been found previously [8]. The full boundary term with the extrinsic curvature and supersymmetry corrections is

$$S_0 = \frac{2}{\kappa^2} \int_{\partial \mathcal{M}} d\mu \left(K - \frac{1}{4} \bar{\psi}_A \Gamma^{AB} \psi_B + \frac{1}{2} \bar{\psi}_A \Gamma^A \psi_N \right) \quad (2)$$

*Electronic address: ian.moss@ncl.ac.uk

Variation of the action $S_{SG} + S_0$ gives a boundary condition on the extrinsic curvature and a chirality condition on the gravitino $P_+ \psi_A = 0$, where $P_+ = (1 + \Gamma_N)/2$. These boundary conditions are consistent with the symmetries of the orbifold S^1/Z_2 . The Bianchi identity implies an additional boundary condition $G_{ABCD} = 0$ which cannot be derived from the action.

The gauge multiplet contains an E_8 gauge field A_A^a and chiral fermions χ^a in the adjoint representation. The action is based on the Yang-Mills action with an interaction term involving the supercurrent,

$$S_1 = -\frac{2\epsilon}{\kappa^2} \int_{\partial\mathcal{M}} d\mu \left(\frac{1}{4} F^a{}_{AB} F^{aAB} + \frac{1}{2} \bar{\chi}^a \Gamma^A D_A(\hat{\Omega}) \chi^a + \frac{1}{4} \bar{\psi}_A \Gamma^{BC} \Gamma^A F^{a*}{}_{BC} \chi^a + \frac{1}{192} \bar{\chi} \Gamma_{ABC} \chi \bar{\psi}_D \Gamma^{ABCDE} \psi_E \right) \quad (3)$$

The constant ϵ sets the relative scale of the matter coupling. The total action $S = S_{SG} + S_0 + S_1$. Note that $F^* = (F + \hat{F})/2$, where

$$\hat{F}^a{}_{AB} = F^a{}_{AB} - \bar{\psi}_{[A} \Gamma_{B]} \chi^a. \quad (4)$$

The complicated four fermi interaction in the action is required for obtaining supercovariant boundary conditions.

One boundary condition is imposed as a constraint. The tangential components of the three form will be required to satisfy $c_{ABC} = 0$, where

$$c_{ABC} = C_{ABC} + \frac{\sqrt{2}}{12} \epsilon \omega_{ABC} + \frac{\sqrt{2}}{48} \epsilon \bar{\chi}^a \Gamma_{ABC} \chi^a \quad (5)$$

This boundary condition contains the Chern-Simons form,

$$\omega = \text{tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right). \quad (6)$$

We shall see shortly how this boundary condition is fixed by supersymmetry, but first it is interesting to check consistency with the gauge symmetry. Under a non-abelian gauge transformation with $\delta A_A^a = -D_A \epsilon^a$, the variation in the Chern-Simons form becomes $\delta \omega = d(\epsilon^a F^a)$. The non-abelian gauge transformation of the Chern-Simons form can therefore be absorbed by an abelian gauge transformation of the three-form (in analogy with Yang-Mills Supergravity [9]). However, the $C \wedge G \wedge G$ term in the supergravity action is not invariant under the abelian gauge transformation, and a quantum gauge anomaly has to be used to restore the gauge symmetry. This is the generalised Green-Schwarz [10] mechanism described by Horava and Witten [2], and it fixes the expansion parameter,

$$\epsilon = \frac{1}{4\pi} \left(\frac{\kappa}{4\pi} \right)^{2/3}. \quad (7)$$

There are also gravitational anomalies which add $R \wedge R$ terms to the boundary condition (5). However, for the present we shall restrict attention to the low curvature regime and ignore $R \wedge R$ terms.

The remaining boundary conditions can be obtained by finding the extrema of the action. Leaving the gauge multiplet fixed for the moment, the relevant boundary terms in the variation of the action are

$$\delta S = \frac{2}{\kappa^2} \int_{\partial\mathcal{M}} d\mu \left(\delta g_{AB} p^{AB} + \delta \bar{\psi}_i \theta^i \right), \quad (8)$$

with

$$p^{AB} = -\frac{1}{2} (K^{AB} - K g^{AB}) + \frac{1}{4} \kappa^2 T^{AB} \quad (9)$$

$$\theta^A = -\Gamma^{AB} P_+ \psi_B - \frac{\epsilon}{4} \Gamma^{BC} \Gamma^A \hat{F}^a{}_{BC} \chi^a - \frac{\epsilon}{96} \Gamma^{AB} \Gamma^{CDE} \psi_B \bar{\chi} \Gamma_{CDE} \chi, \quad (10)$$

where T^{AB} is the surface stress energy tensor.

The boundary condition $p^{AB} = 0$ corresponds, in the covering space, to a junction condition across the brane [7]. The boundary condition $\theta^A = 0$ is a chirality condition on the gravitino

$$P_+ \psi_A = \frac{\epsilon}{12} (\Gamma_A{}^{BC} - 10 \delta_A{}^B \Gamma^C) \hat{F}^a{}_{BC} \chi^a - \frac{\epsilon}{96} \bar{\chi} \Gamma_{BCD} \chi \Gamma^{BCD} P_- \psi_A \quad (11)$$

This boundary condition represents a significant difference between the present model and the construction of Horava and Witten [16]. One significant difference is that torsion can be generated by the matter fields. This affects the connection $\hat{\Omega}$ which governs the motion of gauginos.

The chirality condition on the gravitino and the boundary condition on the three form play a special role in the supersymmetry of the theory. The supersymmetry transformation rules are almost conventional,

$$\delta g_{IJ} = \bar{\eta} \Gamma_{(I} \psi_{J)} \quad (12)$$

$$\delta \psi_I = D_I(\hat{\Omega})\eta + \frac{\sqrt{2}}{288} (\Gamma_I^{JKLM} - 8\delta_I^J \Gamma^{KLM}) \eta \hat{G}_{JKLM} \quad (13)$$

$$\delta C_{IJK} = -\frac{\sqrt{2}}{8} \bar{\eta} \Gamma_{[IJ} \psi_{K]} + \frac{\sqrt{2}}{4} \epsilon \partial_{[I} f_{JK]} \quad (14)$$

$$\delta A^a{}_A = \frac{1}{2} \bar{\eta} \Gamma_A \chi^a \quad (15)$$

$$P_- \delta \chi^a = -\frac{1}{4} \Gamma^{AB} \hat{F}^a{}_{AB} \eta \quad (16)$$

The two-form f will be explained below. The restriction on the chirality of the supersymmetry parameter is less conventional, but is determined by the need to ensure supercovariance of the fermion boundary condition (11),

$$P_+ \eta = -\frac{\epsilon}{96} \bar{\chi} \Gamma_{ABC} \chi \Gamma^{ABC} P_- \eta. \quad (17)$$

This can be regarded as a modification of the projection operator P_+ . The chirality condition on the gaugino is simply $P_+ \chi = 0$ and is not affected by any modification due to a convenient Fierz rearrangement.

We shall consider the supersymmetry of the boundary conditions first, and then move on to the supersymmetry of the action. The variation of C_{ABC} introduces a two-form f , which can be any two-form with the boundary condition

$$f_{AB} = 2A^a{}_{[A} \delta A^a{}_{B]} \quad (18)$$

The two-form is needed to cancel a gauge non-invariant term from the variation of the Chern Simons form in the boundary condition (5). The remaining terms in the variation of C_{ABC} depend only on $P_+ \psi_A$, allowing us to use the gravitino boundary condition to show consistency with the variation of the remaining terms. A calculation has been done to confirm that the boundary condition (5) is supersymmetric to *all* orders in ϵ . A more detailed discussion will be presented in a longer publication.

Taking the exterior derivative of $c_{ABC} = 0$ and making use of (11) puts the boundary condition in manifestly gauge invariant (and supercovariant) form,

$$\hat{G}_{ABCD} = -3\sqrt{2}\epsilon \hat{F}^a{}_{[AB} \hat{F}^a{}_{CD]} + \sqrt{2}\epsilon \bar{\chi}^a \Gamma_{[ABC} D_{D]}(\hat{\Omega})\chi^a + \frac{\sqrt{2}}{4} \epsilon \bar{\chi}^a \Gamma_{[ABC} \Gamma^{EF} \psi_{D]} \hat{F}^a{}_{EF}. \quad (19)$$

This can be used to check the supersymmetry of the gravitino boundary condition (11).

The variation of the action under the supersymmetry transformations can be obtained by combining the boundary terms (8) with terms from the variation of the matter multiplet and with terms which arise from the interior. The invariance of 11-dimensional supergravity ensures that the volume terms in the variation cancel. Boundary terms arise from the interior when partial integration has to be used.

If we write the gravitino variation in the interior in the form $\delta \psi_I L^I$, then integration by parts adds a term $\bar{\eta} L_N$ to the other boundary terms in (8), where

$$L_N = -\Gamma_N \Gamma^{AB} D_A(\hat{\Omega})\psi_B - \frac{\sqrt{2}}{96} \Gamma_N \Gamma^{ABCDE} \psi_A \hat{G}_{BCDE} + \frac{\sqrt{2}}{8} \Gamma_N \Gamma^{AB} \psi^C \hat{G}_{ABCN} \quad (20)$$

A slight rearrangement gives

$$\begin{aligned} \bar{\eta} L_N &= \bar{\eta} D_A(\hat{\Omega})\theta^A + \frac{1}{2} K_{AC} \bar{\eta} \Gamma^{AB} \Gamma^C \psi_B + \frac{1}{4} \epsilon \bar{\eta} D_A(\hat{\Omega}) \left(\Gamma^{BC} \Gamma^A \hat{F}_{BC} \chi \right) \\ &\quad - \frac{\sqrt{2}}{96} \bar{\eta} \Gamma^{ABCDE} \psi_A \hat{G}_{BCDE} + \frac{\sqrt{2}}{8} \bar{\eta} \Gamma^{AB} \psi^C \hat{G}_{ABCN}. \end{aligned} \quad (21)$$

We see clearly the term which cancels the extrinsic curvature in (9) and a term involving the Yang-Mills supercurrent which partially cancels the variation of the Yang-Mills part of the action. The penultimate term cancels the variation of the supercurrent (using 19) and the final term partially cancels with the variation of C_{ABC} .

The three-form and gravitino boundary conditions have to be imposed in order to obtain a supersymmetric action. After some calculation, the action is found to be supersymmetric up to terms of order ϵ^3 (equivalent to κ^2). At this

order there are terms arising from the variation of the $C \wedge G \wedge G$ term in the action and no obvious source of terms for these to cancel. This is an advance over previous attempts to construct the action which have encountered infinite terms at order $\kappa^{4/3}$.

The action and boundary conditions provide a supersymmetric theory which is a natural candidate for the a low energy limit of M -theory. Further work is needed to investigate fully what happens to the supersymmetry at order κ^2 and to complete the supersymmetry variation of the gravitino boundary condition. Going beyond the low energy limit, it would be desirable to include the $R \wedge R$ curvature terms which are significant in compactifications with Calabi-Yau manifolds [11, 12, 13, 14, 15].

-
- [1] P. Horava and E. Witten, Nucl. Phys. **B460**, 506 (1996).
 - [2] P. Horava and E. Witten, Nucl. Phys. **B475**, 94 (1996).
 - [3] E. Bergshoeff, M. de Roo, B. de Ditt, and P. van Nieuwenhuisen, Nucl. Phys. **B195**, 97 (1982).
 - [4] E. Cremmer, B. Julia, and J. Scherk, Phys Lett **B76**, 409 (1978).
 - [5] J. W. York, Phys Rev Lett **28**, 1082 (1972).
 - [6] G. W. Gibbons and S. W. H. Hawking, Phys Rev **D15**, 2752 (1977).
 - [7] H. A. Chamblin and H. S. Reall, Nucl Phys B **133**, 133 (1999).
 - [8] H. C. Luckock and I. G. Moss, Class Quantum grav **6**, 1993 (1989).
 - [9] G. Chapline and N. S. Manton, Phys. Lett **B120**, 105 (1983).
 - [10] M. B. Green and J. H. Schwarz, Phys Lett **B149**, 117 (1984).
 - [11] E. Witten, Nucl. Phys. **B471**, 135 (1996).
 - [12] T. Banks and M. Dine, Nucl Phys **B479**, 173 (1996).
 - [13] A. Lukas, B. A. Ovrut, K. S. Stelle, and D. Waldram, Phys Rev D **59**, 086001 (1999).
 - [14] A. Lukas, B. A. Ovrut, K. S. Stelle, and D. Waldram, Nucl Phys B **552**, 246 (1999).
 - [15] J. Ellis, Z. Lalak, S. Pokorski, and W. pokorski, Nucl Phys **B540**, 149 (1999).
 - [16] The supersymmetry transformation of the chirality condition $\Gamma_{11}\psi_A = \psi_A$ used by Horava and Witten [2] in the covering space approach is ambiguous because the supersymmetry transformation of the gravitino is discontinuous across the brane.